



Scan the QR code to see the full directory of facilities!

Code: 11511

Taking Care of Your Cast



The information provided in this publication is meant purely for educational purposes and may not be used as a substitute for medical diagnosis or treatment. You should seek the advice of your doctor or a qualified healthcare provider before starting any treatment or if you have any questions related to your health, physical fitness or medical conditions. Information is accurate at time of printing (Jan 2023) and subject to revision without prior notice.

^{© 2022.} Alexandra Hospital. All rights reserved. No part of this publication may be reproduced without permission in writing from Alexandra Hospital.

To help your injury heal in the fastest time possible, we have immobilised the injured part of your body with a Plaster-of-Paris (POP) or fiberglass cast. While there may be some initial discomfort, do take heart as you will get used to it and it will be taken off soon.

Feeling tight?

The plaster may feel slightly tight for some time after application.

What are some dos and dont's?

Do's

- Keep the injured hand or wrist above the elbow while lying down.
- Keep the injured foot or knee higher than the hip while lying down.
- Elevate the injured part as long as there is swelling.
- Constantly move the joints not covered by the plaster.
- Keep your cast dry at all times, especially while bathing. When bathing, you may use a plastic sheet to keep your cast dry.

Don'ts

- Do not scratch any itch under the cast as doing so will displace the padding and create a constriction band, which affects blood flow to the injured area.
- Remove the padding under the cast as it serves to provide comfort, protect bony prominences, and prevent friction.

When should I seek medical help?

Consult a doctor at the **Alexandra Hospital Urgent Care Centre (UCC)** if you experience any of the following symptoms:

- If the plaster cast cracks or becomes loose or uncomfortable
- If there is increasing pain or discharge
- If your fingers or toes become swollen, numb or difficult to move
- If there is discolouration of fingers or toes

Notes: